

L. Kaufman, K. Kumar
University of Massachusetts, Amherst, MA 01003

F. W. Hersman
University of New Hampshire, Durham, NH 03824

V. Punjabi
Norfolk State University, Norfolk, VA 23504

L. Todor, L. Weinstein
Old Dominion University, Norfolk, VA 23508

G. Cates, B. Humensky, I. Kominis
Princeton University, Princeton, NJ 08544

E. Brash
University of Regina, Regina, SK, S4S OA2, Canada

S. Dieterich, R. Gilman, C. Glashauser, X. Jiang, G. Kumbartzki,
R. Ransome, S. Strauch
Rutgers University, Piscataway, NJ 08855

L. Auerbach, S. Choi (Spokesperson), Z.-E. Meziani, K. Slifer
Temple University, Philadelphia, PA 19122

H. Tsubota
Tohoku University, Sendai, Japan

D. Armstrong, D. T. Averett, P. Djawotho, M. Finn, K. Griffioen, K. Kramer
College of William and Mary, Williamsburg, VA 23185

June 1, 2000

Abstract

We propose a precision extraction of the neutron spin structure function g_1^n and the virtual photon asymmetry A_1^n in the resonance region over a moderate Q^2 range (up to $Q^2 = 4.2(\text{GeV}/c)^2$) using a polarized ${}^3\text{He}$ target. The Bloom-Gilman duality has been experimentally demonstrated for the spin independent structure functions down to small values of Q^2 . The proposed experiment combined with Deep-Inelastic-Scattering data will provide a precision test of quark-hadron duality predictions for spin structure functions. The demonstration of duality for spin structure functions will enable the use of resonance data as a powerful tool to study the nucleon spin structure in the very high x region.